



# BRADYCARDIA WITH PULSE



## Escambia County, Florida - ALS/BLS Medical Protocol

---

### Supportive Care

1. Maintain airway, Oxygen if hypoxemic
2. EKG monitor, O2 sat. monitor, obtain IV access
3. 12 lead EKG, transmit to receiving hospital (don't delay treatment)
4. Follow [Medical Supportive Care Protocol](#).

### ALS Level 1


1. **IF UNSTABLE WITH** signs or symptoms of HYPOTENSION, ALTERED MENTAL STATUS, SIGNS OF SHOCK, ISCHEMIC CHEST PAIN, or ACUTE HEART FAILURE then
  - 1a. [Atropine](#) 0.5mg bolus IV (First Dose). Repeat every 3 -5 minutes. Total Max. dose = 3mg.  
  
or
  - 1b. If atropine ineffective begin [transcutaneous pacing](#)  
  
or
  - 1c. [Dopamine](#) IV infusion 2-10 mcg/kg/min.  
  
or
  - 1d. [Epinephrine](#) IV infusion 2-10 mcg/min.
2. **IF STABLE WITHOUT** signs or symptoms of HYPOTENSION, ALTERED MENTAL STATUS, SIGNS OF SHOCK, ISCHEMIC CHEST PAIN, or ACUTE HEART FAILURE) then
  - 2a. Monitor and observe



Approved: 10/01/2011  
Version: 3.1

Page 1 of 1

Approved by:

  
Charles Neal, D.O. Medical Director