# PEDIATRIC BRADYCARDIA

**Escambia County, Florida - ALS/BLS Medical Protocol**

## Supportive Care

1. Maintain airway, Oxygen if hypoxemic
2. EKG monitor, O2 sat. monitor, obtain IV access
3. 12 lead EKG, transmit to receiving hospital (don't delay treatment)
4. Follow Medical Supportive Care Protocol.

## ALS Level 1

**1.** **IF UNSTABLE *WITH*** signs or symptoms of HYPOTENSION, ALTERED MENTAL STATUS, SIGNS OF SHOCK**,** ISCHEMIC CHEST PAIN, or ACUTE HEART FAILURE then

1a. CPR if heart rate < 60 per minute.

1b. **Epinephrine** IV 0.01 mg/kg.

1c. **Atropine** 0.02 mg/kg bolus IV.

May repeat once.

Minimum single dose 0.1 mg, Maximum single dose 0.5 mg.

1d. If epinephrine and atropine ineffective begin transcutanous pacing.

**2.** **IF STABLE *WITHOUT*** signs or symptoms of HYPOTENSION, ALTERED MENTAL STATUS, SIGNS OF SHOCK, ISCHEMIC CHEST PAIN, or ACUTE HEART FAILURE) then

2a. Monitor and observe